


A probabilistic reinforcement-learning algorithm to find shortest paths in a graph

Zoé Varin (IRIF)

January 20th, 2026

Joint work with Cécile Mailler (University of Bath)



- Since November: Postdoc in the Combinatorics team
 - working with Marie Albenque and Guillaume Chapuy
 - funded by the ComplexCité consortium
- PhD in Bordeaux (LaBRI) supervised by Jean-François Marckert
 - parking model and various generalisations
 - [the ant model \(with Cécile Mailler\)](#)

Introduction



Two biological ants' experiments:




FIGURE 1 Experimental setup for the double bridge experiment.
(a) Branches have equal lengths [3]. (b) Branches have different lengths [4].

Two biological ants' experiments:




FIGURE 1 Experimental setup for the double bridge experiment. (a) Branches have equal lengths [3]. (b) Branches have different lengths [4].

Models to fit the experimental results:

$$p_1 = \frac{(m_1 + k)^h}{(m_1 + k)^h + (m_2 + k)^h}, \quad (1)$$

(p_1 = probability to go in branch 1; m_i = quantity of pheromones in branch i ; k and h parameters to fitted)

Two biological ants' experiments:




FIGURE 1 Experimental setup for the double bridge experiment. (a) Branches have equal lengths [3]. (b) Branches have different lengths [4].

Models to fit the experimental results:


$$p_1 = \frac{(m_1 + k)^h}{(m_1 + k)^h + (m_2 + k)^h}, \quad (1)$$

(p_1 = probability to go in branch 1; m_i = quantity of pheromones in branch i ; k and h parameters to fitted)

→ **Definition of a Metaheuristic: Ant Colony Optimization**

📖 See *Ant Colony Optimization* (Dorigo, Birattari and Stützle) for a nice introduction (and the source of the screenshots)

Definition of the model (one-nest version)




At each step n :

- **random walk** X , starting from \mathcal{N} , stopped at \mathcal{F}


Definition of the model (one-nest version)

At each step n :

- **random walk** X , starting from \mathcal{N} , stopped at \mathcal{F}



Definition of the model (one-nest version)




At each step n :

- **random walk** X , starting from \mathcal{N} , stopped at \mathcal{F}

- **depositing pheromones** on γ on their way back. Loop-erased (LE) model: $\gamma = LE(X)$

Definition of the model (one-nest version)



At each step n :


- **random walk** X , starting from \mathcal{N} , stopped at \mathcal{F}

- **depositing pheromones** on γ on their way back. Loop-erased (LE) model: $\gamma = LE(X)$

- **Edge weights:**

$$W_e(n) = 1 + \#\text{pheromones on } e$$

Definition of the model (one-nest version)



At each step n :

- **random walk** X , starting from \mathcal{N} , stopped at \mathcal{F} and weighted by $W(n)$:


$$\mathbb{P}(u \rightarrow v) = \frac{W_{uv}(n)}{\sum_{e:u \in e} W_e(n)}$$

- **depositing pheromones** on γ on their way back. Loop-erased (LE) model: $\gamma = LE(X)$

- **Edge weights:**

$$W_e(n) = 1 + \#\text{pheromones on } e$$

Definition of the model (one-nest version)



At each step n :


- **random walk** X , starting from \mathcal{N} , stopped at \mathcal{F} and weighted by $W(n)$:

$$\mathbb{P}(u \rightarrow v) = \frac{W_{uv}(n)}{\sum_{e:u \in e} W_e(n)}$$

- **depositing pheromones** on γ on their way back. Loop-erased (LE) model: $\gamma = LE(X)$


- **Edge weights:**

$$W_e(n) = 1 + \#\text{pheromones on } e$$



Simulations for $n = 10^8$

Definition of the model (one-nest version)



At each step n :


- **random walk** X , starting from \mathcal{N} , stopped at \mathcal{F} and weighted by $W(n)$:

$$\mathbb{P}(u \rightarrow v) = \frac{W_{uv}(n)}{\sum_{e:u \in e} W_e(n)}$$

- **depositing pheromones** on γ on their way back. Loop-erased (LE) model: $\gamma = LE(X)$


- **Edge weights:**

$$W_e(n) = 1 + \#\text{pheromones on } e$$



Simulations for $n = 10^8$

Definition of the model (one-nest version)



At each step n :

- **random walk** X , starting from \mathcal{N} , stopped at \mathcal{F} and weighted by $W(n)$:

$$\mathbb{P}(u \rightarrow v) = \frac{W_{uv}(n)}{\sum_{e:u \in e} W_e(n)}$$

- **depositing pheromones** on γ on their way back. Loop-erased (LE) model: $\gamma = LE(X)$


- **Edge weights:**

$$W_e(n) = 1 + \#\text{pheromones on } e$$

Question: Do the ants find shortest paths from \mathcal{N} to \mathcal{F} ?

→ Does $\left(\frac{W_e(n)}{n}\right)_e$ converge ?

Towards which limit ?




Simulations for $n = 10^8$

A quick warm-up




Pólya's urn:




A quick warm-up




Pólya's urn:



A quick warm-up




Pólya's urn:



A quick warm-up




Pólya's urn:



A quick warm-up




Pólya's urn:



A quick warm-up




Pólya's urn:



A quick warm-up




Pólya's urn:



A quick warm-up




Pólya's urn:



A quick warm-up




Pólya's urn:



A quick warm-up




Pólya's urn:



A quick warm-up




Pólya's urn:



A quick warm-up



Pólya's urn:



A quick warm-up



Pólya's urn:



A quick warm-up



Pólya's urn:




Asymptotic behavior:

Almost surely,

$$\frac{\#\text{orange}}{n} \xrightarrow[n \rightarrow \infty]{} U \sim \mathcal{U}([0, 1])$$


Loop-erased (LE) model on series-parallel graphs

Recursive definition of series-parallel (SP) graphs:



Loop-erased (LE) model on series-parallel graphs

Recursive definition of series-parallel (SP) graphs:



Theorem (Kious, Mailler, Schapira 22)


If G is a SP graph, then almost surely,

$$\frac{W_e(n)}{n} \xrightarrow[n \rightarrow \infty]{} \chi_e, \quad \forall e \in E$$

where $(\chi_e)_{e \in E}$ is a random vector such that $\forall e, \chi_e \neq 0 \iff e \in \text{Geodesic}(G)$.

Loop-erased (LE) model on series-parallel graphs

Recursive definition of series-parallel (SP) graphs:




Theorem (Kious, Mailler, Schapira 22)


If G is a SP graph, then almost surely,

$$\frac{W_e(n)}{n} \xrightarrow[n \rightarrow \infty]{} \chi_e, \quad \forall e \in E$$

where $(\chi_e)_{e \in E}$ is a random vector such that $\forall e, \chi_e \neq 0 \iff e \in \text{Geodesic}(G)$.




Other reinforcement models




(T) trace:

$$\gamma = X$$



(LE) loop-erased:

$$\gamma = LE(X)$$



(G) geodesic:

$$\gamma = \text{ShortestPath}(X)$$

At each step n :


• **random walk** X :

• **depositing pheromones** on γ :

$$\forall e, W_e(n+1) = W_e(n) + \mathbb{1}_{e \in \gamma}$$

Geodesic (G) model on the lozenge graph

The lozenge graph:



Theorem (Kious, Mailler, Schapira 22)

Almost surely,

$$\frac{W_i(n)}{n} \xrightarrow[n \rightarrow \infty]{} \chi_i, \quad \forall i \in \{a, b, c, d, e\}$$

where $(\chi_i)_{i \in \{a, b, c, d, e\}}$ is a random vector, such that almost surely, $\chi_a = \chi_b = 1 - \chi_d = 1 - \chi_e \in (0, 1)$ and $\chi_c = 0$.

Conjecture for the loop-erased (LE) and geodesic (G) models

Conjecture (KMS22)

Almost surely,

$$\frac{W_e(n)}{n} \xrightarrow[n \rightarrow \infty]{} \chi_e, \quad \forall e \in E$$

where $(\chi_e)_{e \in E}$ is a random vector such that

(LE) model $\chi_e \neq 0$ a.s. **if and only if** e belongs to a shortest path from N to F

(G) model $\chi_e \neq 0$ a.s. **only if** e belongs to a shortest path from N to F

Conjecture for the loop-erased (LE) and geodesic (G) models

Conjecture (KMS22)


Almost surely,

$$\frac{W_e(n)}{n} \xrightarrow[n \rightarrow \infty]{} \chi_e, \quad \forall e \in E$$

where $(\chi_e)_{e \in E}$ is a random vector such that

(LE) model $\chi_e \neq 0$ a.s. **if and only if** e belongs to a shortest path from N to F

(G) model $\chi_e \neq 0$ a.s. **only if** e belongs to a shortest path from N to F




For L large enough, there exists e such that


$$\mathbb{P}(W_e(n)/n \rightarrow 0) > 0$$

Trace (T) model (Results from KMS22)

Some examples: the cone and the lozenge



$$\frac{W(n)}{n} \xrightarrow[n \rightarrow \infty]{} (1, 1/3, 1/3, 0)$$



$$\frac{W(n)}{n} \xrightarrow[n \rightarrow \infty]{} (w^*, 1/2, 1/2, w^*, 1/2)$$

Theorem: convergence for a family of *tree-like* graphs (s.t. $G \setminus \{\mathcal{F}\}$ is a tree).


Conjecture: deterministic limit for any graph without multiple-edges adjacent to \mathcal{F} .

The two-nest model




Back to the (LE) model: multinest version

2-nest version: at every step n , $\mathcal{N}(n) = \begin{cases} \mathcal{N}_1 & \text{with proba } \alpha \in (0, 1) \\ \mathcal{N}_2 & \text{with proba } 1 - \alpha \end{cases}$.



Back to the (LE) model: multinest version on triangle-SP graphs

2-nest version: at every step n , $\mathcal{N}(n) = \begin{cases} \mathcal{N}_1 & \text{with proba } \alpha \in (0, 1) \\ \mathcal{N}_2 & \text{with proba } 1 - \alpha \end{cases}$.




Triangle-SP graph: G_1, G_2, G_3 series-parallel graphs

Our main result: convergence of the 2-nest LE model on triangle-SP graphs

$N_i(n)$ = number of steps at which edges in G_i have been reinforced

Triangle-SP graph



Remark: $\forall n, N_1(n) + N_2(n) = n$.

Our main result: convergence of the 2-nest LE model on triangle-SP graphs


$N_i(n)$ = number of steps at which edges in G_i have been reinforced

Theorem (Mailler, V. 25+)

Almost surely,

$$\left(\frac{N_1(n)}{n}, \frac{N_2(n)}{n}, \frac{N_3(n)}{n} \right) \xrightarrow{n \rightarrow \infty} w$$

Triangle-SP graph



Remark: $\forall n, N_1(n) + N_2(n) = n$.

Our main result: convergence of the 2-nest LE model on triangle-SP graphs

$N_i(n)$ = number of steps at which edges in G_i have been reinforced

Theorem (Mailler, V. 25+)


Almost surely,

$$\left(\frac{N_1(n)}{n}, \frac{N_2(n)}{n}, \frac{N_3(n)}{n} \right) \xrightarrow{n \rightarrow \infty} w$$

If $\ell_1 \leq \ell_2$, then,

- if $\ell_2 \geq \ell_1 + \ell_3$, then $w = (1, 0, 1 - \alpha)$,

Triangle-SP graph



Remark: $\forall n, N_1(n) + N_2(n) = n$.

Our main result: convergence of the 2-nest LE model on triangle-SP graphs

$N_i(n)$ = number of steps at which edges in G_i have been reinforced

Theorem (Mailler, V. 25+)


Almost surely,

$$\left(\frac{N_1(n)}{n}, \frac{N_2(n)}{n}, \frac{N_3(n)}{n} \right) \xrightarrow{n \rightarrow \infty} w$$

If $\ell_1 \leq \ell_2$, then,

- if $\ell_2 \geq \ell_1 + \ell_3$, then $w = (1, 0, 1 - \alpha)$,
- if $\ell_3 \geq \ell_1 + \ell_2$, then $w = (\alpha, 1 - \alpha, 0)$,

Triangle-SP graph



Remark: $\forall n, N_1(n) + N_2(n) = n$.

Our main result: convergence of the 2-nest LE model on triangle-SP graphs

$N_i(n)$ = number of steps at which edges in G_i have been reinforced

Theorem (Mailler, V. 25+)

Almost surely,

$$\left(\frac{N_1(n)}{n}, \frac{N_2(n)}{n}, \frac{N_3(n)}{n} \right) \xrightarrow{n \rightarrow \infty} w$$

If $l_1 \leq l_2$, then,

- if $l_2 \geq l_1 + l_3$, then $w = (1, 0, 1 - \alpha)$,
- if $l_3 \geq l_1 + l_2$, then $w = (\alpha, 1 - \alpha, 0)$,
- otherwise $w = (\beta_1, 1 - \beta_1, \beta_3)$.

$$\beta_1 = \frac{\alpha_1 l_1 (l_3 + l_2 - l_1)}{l_1 l_3 + (l_2 - l_1) ((1 - \alpha_1)(l_3 - l_2) + \alpha_1 l_1)}$$

$$\beta_3 = \frac{\alpha_1 l_3 (1 - \alpha_1) (l_1 + l_2 - l_3)}{(l_2 - l_1) (l_1 + l_2 - l_3) \alpha_1 + l_2 (l_1 - l_2 + l_3)}$$

$= d_{G_2}(\mathcal{N}_2, \mathcal{F})$

Remark: $\forall n, N_1(n) + N_2(n) = n$.



Our main result: convergence of the 2-nest LE model on triangle-SP graphs

$N_i(n)$ = number of steps at which edges in G_i have been reinforced

Theorem (Mailler, V. 25+)

Almost surely,

$$\left(\frac{N_1(n)}{n}, \frac{N_2(n)}{n}, \frac{N_3(n)}{n} \right) \xrightarrow{n \rightarrow \infty} w$$


If $\ell_1 \leq \ell_2$, then,

- if $\ell_2 \geq \ell_1 + \ell_3$, then $w = (1, 0, 1 - \alpha)$,
- if $\ell_3 \geq \ell_1 + \ell_2$, then $w = (\alpha, 1 - \alpha, 0)$,
- otherwise $w = (\beta_1, 1 - \beta_1, \beta_3)$.

Almost surely, $\forall e \in G_i, \frac{W_e(n)}{n} \xrightarrow{n \rightarrow \infty} \xi_e$ (random),

$$\text{where } \xi_e \neq 0 \iff \begin{cases} e \in \text{Geodesic}(G_i) \\ \lim N_i(n)/n > 0 \end{cases}$$

Triangle-SP graph



Remark: $\forall n, N_1(n) + N_2(n) = n$.

Our main result: convergence of the 2-nest LE model on triangle-SP graphs

$N_i(n)$ = number of steps at which edges in G_i have been reinforced

Theorem (Mailler, V. 25+)

Almost surely,

$$\left(\frac{N_1(n)}{n}, \frac{N_2(n)}{n}, \frac{N_3(n)}{n} \right) \xrightarrow{n \rightarrow \infty} w$$


If $\ell_1 \leq \ell_2$, then,

- if $\ell_2 \geq \ell_1 + \ell_3$, then $w = (1, 0, 1 - \alpha)$,
- if $\ell_3 \geq \ell_1 + \ell_2$, then $w = (\alpha, 1 - \alpha, 0)$,
- otherwise $w = (\beta_1, 1 - \beta_1, \beta_3)$.

Almost surely, $\forall e \in G_i, \frac{W_e(n)}{n} \xrightarrow{n \rightarrow \infty} \xi_e$ (random),

$$\text{where } \xi_e \neq 0 \iff \begin{cases} e \in \text{Geodesic}(G_i) \\ \lim N_i(n)/n > 0 \end{cases}$$

Triangle-SP graph



Our main result: convergence of the 2-nest LE model on triangle-SP graphs

$N_i(n)$ = number of steps at which edges in G_i have been reinforced

Theorem (Mailler, V. 25+)

Almost surely,

$$\left(\frac{N_1(n)}{n}, \frac{N_2(n)}{n}, \frac{N_3(n)}{n} \right) \xrightarrow{n \rightarrow \infty} w$$


If $\ell_1 \leq \ell_2$, then,

- if $\ell_2 \geq \ell_1 + \ell_3$, then $w = (1, 0, 1 - \alpha)$,
- if $\ell_3 \geq \ell_1 + \ell_2$, then $w = (\alpha, 1 - \alpha, 0)$,
- otherwise $w = (\beta_1, 1 - \beta_1, \beta_3)$.

Almost surely, $\forall e \in G_i, \frac{W_e(n)}{n} \xrightarrow{n \rightarrow \infty} \xi_e$ (random),

$$\text{where } \xi_e \neq 0 \iff \begin{cases} e \in \text{Geodesic}(G_i) \\ \lim N_i(n)/n > 0 \end{cases}$$

Triangle-SP graph



When $\ell_1 = \ell_2 = \ell_3$, $\beta_1 = \alpha$ and $\beta_3 = \alpha(1 - \alpha)$.

Our main result: convergence of the 2-nest LE model on triangle-SP graphs

$N_i(n)$ = number of steps at which edges in G_i have been reinforced

Theorem (Mailler, V. 25+)

Almost surely,

$$\left(\frac{N_1(n)}{n}, \frac{N_2(n)}{n}, \frac{N_3(n)}{n} \right) \xrightarrow{n \rightarrow \infty} w$$


If $\ell_1 \leq \ell_2$, then,

- if $\ell_2 \geq \ell_1 + \ell_3$, then $w = (1, 0, 1 - \alpha)$,
- if $\ell_3 \geq \ell_1 + \ell_2$, then $w = (\alpha, 1 - \alpha, 0)$,
- otherwise $w = (\beta_1, 1 - \beta_1, \beta_3)$.

Almost surely, $\forall e \in G_i$, $\frac{W_e(n)}{n} \xrightarrow{n \rightarrow \infty} \xi_e$ (random),

$$\text{where } \xi_e \neq 0 \iff \begin{cases} e \in \text{Geodesic}(G_i) \\ \lim N_i(n)/n > 0 \end{cases}$$

Triangle-SP graph



Toolbox:


- Pólya urns
- Stochastic approximation theory
- Conductance method and results on the one-nest model on SP-graphs

Toolbox & Proof





Conductance method






Conductance method

 See *Probability on trees and networks* (Lyons, Peres 16)

Effective conductance between two vertices - **recursive definition for SP graphs**:




(a) $C_G = w$



(b) $C_G = C_G + C_{\tilde{G}}$



(c) $C_G = \frac{1}{1/C_G + 1/C_{\tilde{G}}}$






Conductance method

📖 See *Probability on trees and networks* (Lyons, Peres 16)

Effective conductance between two vertices - **recursive definition for SP graphs**:




(a) $C_G = w$



(b) $C_G = C_G + C_{\tilde{G}}$



(c) $C_G = \frac{1}{1/C_G + 1/C_{\tilde{G}}}$



💡 **Key idea:** the probability that a random walk starting from S hits T before \tilde{T} is $\frac{C_G}{C_G + C_{\tilde{G}}}$.






Conductance method

See *Probability on trees and networks* (Lyons, Peres 16)

Effective conductance between two vertices - **recursive definition for SP graphs**:



(a) $C_G = w$



(b) $C_G = C_G + C_{\tilde{G}}$




(c) $C_G = \frac{1}{1/C_G + 1/C_{\tilde{G}}}$

Key idea: the probability that a random walk starting from S hits T before \tilde{T} is $\frac{C_G}{C_G + C_{\tilde{G}}}$.



Example: If the ant starts from \mathcal{N}_1 , the probability to reinforce in G_1 is

$$\frac{C_{G_1}(n)}{C_{G_1}(n) + \frac{C_{G_2}(n)C_{G_3}(n)}{C_{G_2}(n) + C_{G_3}(n)}}$$






Conductance method

See *Probability on trees and networks* (Lyons, Peres 16)

Effective conductance between two vertices - **recursive definition for SP graphs**:



(a) $C_G = w$



(b) $C_G = C_G + C_{\tilde{G}}$




(c) $C_G = \frac{1}{1/C_G + 1/C_{\tilde{G}}}$

Key idea: the probability that a random walk starting from S hits T before \tilde{T} is $\frac{C_G}{C_G + C_{\tilde{G}}}$.



Example: If the ant starts from \mathcal{N}_1 , the probability to reinforce in G_1 is

$$\frac{C_{G_1}(n)}{C_{G_1}(n) + \frac{C_{G_2}(n)C_{G_3}(n)}{C_{G_2}(n) + C_{G_3}(n)}}$$



Asymptotics for the conductances

On SP graphs:



$$h_{\min}(G) = d_G(\mathcal{N}, \mathcal{F})$$

Theorem (Kious, Mailler, Schapira 22): For every $i \in \{1, 2, 3\}$,

$$\frac{C_G(n)}{n} \xrightarrow{n \rightarrow \infty} \frac{1}{h_{\min}(G)}$$

Asymptotics for the conductances

On SP graphs:



$$h_{\min}(G) = d_G(\mathcal{N}, \mathcal{F})$$

Theorem (Kious, Mailler, Schapira 22): For every $i \in \{1, 2, 3\}$,

$$\frac{C_G(n)}{n} \xrightarrow{n \rightarrow \infty} \frac{1}{h_{\min}(G)}$$

💡 Combinatorial analysis of LE walks

Asymptotics for the conductances

On SP graphs:




$$h_{\min}(G) = d_G(\mathcal{N}, \mathcal{F})$$

Theorem (Kious, Mailler, Schapira 22): For every $i \in \{1, 2, 3\}$,

$$\frac{C_G(n)}{n} \xrightarrow{n \rightarrow \infty} \frac{1}{h_{\min}(G)}$$

 Combinatorial analysis of LE walks



Proposition (Mailler, V. 25+)

For every $i \in \{1, 2, 3\}$,

$$\frac{C_{G_i}(n)}{N_i(n)} \xrightarrow{n \rightarrow \infty} \frac{1}{h_{\min}(G_i)} = \frac{1}{l_i}$$

(Partial) summary of the proof

Step 1: Conductance method. $\mathbb{P}(N_1(n+1) - N_1(n) = 1) = \text{Function}(C_{G_1}, C_{G_2}, C_{G_3})$.

Step 2: $C_{G_i}(n) \underset{n \rightarrow \infty}{\sim} \frac{N_i(n)}{\ell_i}$. (KMS22 + combinatorial analysis of the walks)

(Partial) summary of the proof

Step 1: Conductance method. $\mathbb{P}(N_1(n+1) - N_1(n) = 1) = \text{Function}(C_{G_1}, C_{G_2}, C_{G_3})$.

Step 2: $C_{G_i}(n) \underset{n \rightarrow \infty}{\sim} \frac{N_i(n)}{\ell_i}$. (KMS22 + combinatorial analysis of the walks)

Step 3: Stochastic approximation. The error made in Step 2 is asymptotically negligible.
Thus $\left(\frac{N_1(n)}{n}, \frac{N_3(n)}{n}\right)_{n \geq 0}$ is a stochastic approximation + converges.



Stochastic approximation

📖 See *Random processes with reinforcement* (Pemantle 07)

A process $(X_n)_{n \geq 0}$ is a **stochastic approximation** if

$$X_{n+1} - X_n = \frac{F(X_n) + \xi_{n+1} + r_n}{n+1}, \forall n$$



Stochastic approximation

 See *Random processes with reinforcement* (Pemantle 07)

A process $(X_n)_{n \geq 0}$ is a **stochastic approximation** if

$$X_{n+1} - X_n = \frac{F(X_n) + \xi_{n+1} + r_n}{n+1}, \quad \forall n$$

and if

- $F : \mathcal{E} \rightarrow \mathbb{R}^d$ is a Lipschitz function,
- the noise ξ_{n+1} is \mathcal{F}_{n+1} -measurable and such that $\forall n, \mathbb{E}_n [\xi_{n+1}] = 0$,
- the remainder term r_n is \mathcal{F}_n -measurable and such that $\sum_n n^{-1} \|r_n\| < \infty$ a.s.



Stochastic approximation

📖 See *Random processes with reinforcement* (Pemantle 07)

A process $(X_n)_{n \geq 0}$ is a **stochastic approximation** if

$$X_{n+1} - X_n = \frac{F(X_n) + \xi_{n+1} + r_n}{n+1}, \quad \forall n$$

and if

- $F : \mathcal{E} \rightarrow \mathbb{R}^d$ is a Lipschitz function,
- the noise ξ_{n+1} is \mathcal{F}_{n+1} -measurable and such that $\forall n, \mathbb{E}_n [\xi_{n+1}] = 0$,
- the remainder term r_n is \mathcal{F}_n -measurable and such that $\sum_n n^{-1} \|r_n\| < \infty$ a.s.


Claim: the process $\left(\frac{N_1(n)}{n}, \frac{N_3(n)}{n} \right)_{n \geq 0}$ is a stochastic approximation !

The ODE method

A process $(X_n)_{n \geq 0}$ is a **stochastic approximation** if

$$X_{n+1} - X_n = \frac{F(X_n) + \xi_{n+1} + r_n}{n+1}, \quad \forall n$$

Vector field $F(w_1, w_3)$:




(example with $\ell_1 = 2$, $\ell_2 = 4$ and $\ell_3 = 3$)

The ODE method

A process $(X_n)_{n \geq 0}$ is a **stochastic approximation** if

$$X_{n+1} - X_n = \frac{F(X_n) + \xi_{n+1} + r_n}{n+1}, \quad \forall n$$

Vector field $F(w_1, w_3)$:



(example with $\ell_1 = 2$, $\ell_2 = 4$ and $\ell_3 = 3$)

💡 **Main idea:** if ξ_{n+1} and r_n behave nicely, $(X_n)_{n \leq 0}$ follows the flow of the ODE $\dot{y} = F(y)$!

ODE method

If any solution of the ODE $\dot{y} = F(y)$ converges, then almost surely,

$$\exists \bullet : X_n \xrightarrow{n \rightarrow \infty} \bullet$$


where \bullet is a (random) 0 of F .

The ODE method

A process $(X_n)_{n \geq 0}$ is a **stochastic approximation** if


$$X_{n+1} - X_n = \frac{F(X_n) + \xi_{n+1} + r_n}{n+1}, \forall n$$

Vector field $F(w_1, w_3)$:



(example with $\ell_1 = 2$, $\ell_2 = 4$ and $\ell_3 = 3$)

What does not happen:



Conclusion: any solution to $\dot{y} = F(y)$ starting in $[0, 1]^2$ converges
 $\rightarrow \left(\frac{N_1(n)}{n}, \frac{N_3(n)}{n} \right)$ converges a.s. !

Conclusion of the proof

Step 1: Conductance method. $\mathbb{P}(N_1(n+1) - N_1(n) = 1) = \text{Function}(C_{G_1}, C_{G_2}, C_{G_3})$.

Step 2: $C_{G_i}(n) \underset{n \rightarrow \infty}{\sim} \frac{N_i(n)}{\ell_i}$. (KMS22 + combinatorial analysis of the walks)

Step 3: Stochastic approximation. The error made in Step 2 is asymptotically negligible.

Thus $\left(\frac{N_1(n)}{n}, \frac{N_3(n)}{n}\right)_{n \geq 0}$ is a stochastic approximation + converges.

Conclusion of the proof

Step 1: Conductance method. $\mathbb{P}(N_1(n+1) - N_1(n) = 1) = \text{Function}(C_{G_1}, C_{G_2}, C_{G_3})$.

Step 2: $C_{G_i}(n) \underset{n \rightarrow \infty}{\sim} \frac{N_i(n)}{\ell_i}$. (KMS22 + combinatorial analysis of the walks)

Step 3: Stochastic approximation. The error made in Step 2 is asymptotically negligible. Thus $\left(\frac{N_1(n)}{n}, \frac{N_3(n)}{n}\right)_{n \geq 0}$ is a stochastic approximation + converges.

Step 4: Comparison with generalized Pólya urns to determine the limit.

A lot of unanswered questions remain

- Can we characterize the distribution of the edge weights' limit ?
- What happens on other families of graphs ?
- Or if we consider graphs with more nests ?
- Evaporation of the pheromones ?

Thank you !

